

TIFR/TH/98-09

August, 1998

hep-ph/9803424

# $\eta_c$ Production at the Tevatron: A test of NRQCD

Prakash Mathews<sup>1</sup>, P. Poulose<sup>2</sup> and K. Sridhar<sup>3</sup>

*Department of Theoretical Physics  
Tata Institute of Fundamental Research  
Homi Bhabha Road, Bombay 400005, INDIA.*

## Abstract

We use NRQCD to predict the cross section for  $\eta_c$  production at the Tevatron. The non-perturbative matrix elements required are obtained, using heavy-quark symmetry, from the matrix elements determined from the CDF  $J/\psi$  data. Our numbers are, therefore, predictions of NRQCD and provide a very good test of it. Even after taking into account the small branching ratio of the  $\eta_c$  into two photons, a substantial rate for the production of this resonance at the Tevatron is expected.

(To appear in Physics Letters B)

---

<sup>1</sup>E-mail: prakash@theory.tifr.res.in

<sup>2</sup>E-mail: poulose@theory.tifr.res.in

<sup>3</sup>E-mail: sridhar@theory.tifr.res.in

It is by now well known from compelling theoretical [1] and phenomenological [2] reasons that the Colour Singlet model (CSM) used to describe the production and decay of quarkonium is incomplete. A new factorisation approach within QCD was provided by Bodwin, Braaten and Lepage [3]. This method separates the perturbatively calculable short-distance ( $\leq 1/M_Q$ ,  $M_Q$  is the mass of the heavy quark) effects from the long-distance effects, which are described by non-perturbative matrix elements in an effective field theory called non-relativistic QCD (NRQCD). Unlike CSM where factorisation is explicitly violated, factorisation can be shown to be valid in NRQCD, because in this approach the bound state problem is treated in a systematic expansion in powers of  $v$ , the relative velocity between the quarks in the bound state. Consequently, in NRQCD the production and decay of quarkonium takes place via intermediate states in which the  $Q\bar{Q}$  pair has quantum numbers different from those of the physical quarkonium. The  $Q\bar{Q}$  pair in the intermediate state denoted as  $Q\bar{Q}[^{2S+1}L_J^{[1,8]}]$  could be in the colour singlet or octet state and could form the physical quarkonium state by making chromo-electric or -magnetic transitions. Thus the cross section for production of a quarkonium state  $H$  can be written as

$$\sigma(H) = \sum_n \frac{F_n}{M_Q^{d_n-4}} \langle 0 | \mathcal{O}_n^H | 0 \rangle. \quad (1)$$

The coefficients  $F_n$  correspond to the production of  $Q\bar{Q}$  in the angular momentum and colour state denoted by  $n$  and is calculated using perturbative QCD. The non-perturbative part,  $\langle \mathcal{O}_n^H \rangle$  of mass dimension  $d_n$  in NRQCD has a well-defined operator definition and is universal. These matrix elements can be extracted from any one process and then be used to predict other processes where the same matrix elements appear. Though the summation involves an infinite number of terms, the relative magnitude of the various terms is predicted by NRQCD as these matrix elements scale as powers of  $v$ . However, this does not necessarily imply that effects from higher orders in  $v$  will always be small in physical processes, because any observable, like the decay width or the cross section is given by a double expansion in the strong coupling constant  $\alpha_s(M_Q)$  and the relative velocity  $v$ . Indeed, the higher order colour-octet components have important effects in certain processes: for example, in  $\chi_c$  decays into light hadrons, where these octet components are necessary to obtain an infrared safe expression for the decay-width [3], and in large- $p_T$   $J/\psi$  and  $\psi'$  production from the CDF experiment [2] at Tevatron where discrepancies larger than one order of magnitude *vis a vis* CSM predictions could be explained by including the octet components [4, 5].

The non-perturbative matrix elements in NRQCD, are not calculable and have to be obtained by fitting to available data. The matrix elements of the colour-singlet operators are related to the radial wave-functions and can be obtained from decay

widths or lattice calculations, for example, but the colour-octet matrix elements are not known and have only been obtained by fitting NRQCD predictions to the CDF data [6, 7]. Given that the normalisation of the cross sections are not predictions of the theory, it is important to look for other independent quantitative tests of NRQCD. One crucial test of NRQCD was thought to be the production of large- $p_T$  *inelastic*  $J/\psi$  production in  $ep$  collisions at HERA. The inelastic cross section has been measured by both H1 and ZEUS [8]. The inelasticity of the events is ensured by selecting events with  $z < 1$ , where  $z \equiv p_p \cdot p_{J/\psi} / p_p \cdot p_\gamma$ . In addition, a  $p_T$  cut of 1 GeV is used to select the events. Using the values of the non-perturbative inputs obtained from the Tevatron, the theoretical predictions have been compared [9] to these inelastic data. However, there seems to be a problem in predicting the  $z$  distribution because while the colour-singlet cross section dominates in most of the low- $z$  region, the colour-octet contribution grows anomalously in the large- $z$  ( $0.8 < z < 0.9$ ) region and this rise is not seen in the data. However, it is premature to conclude that this is a failure of NRQCD, because at  $z$  values of 0.8 other effects like soft-gluons [10],  $k_T$ -smearing [11] could become important, or even worse, the NRQCD factorisation may break down [12]. The safer conclusion is that the inelastic  $J/\psi$  production process at HERA is not a clean test of NRQCD.

In fact, NRQCD is best tested at Tevatron itself in other large- $p_T$  observables. The polarisation of the produced  $J/\psi$  is one such test [13]. The second test is to look for the production of other charmonium states at large- $p_T$  at the Tevatron. This is the philosophy of the present paper and we study large- $p_T$   $\eta_c$  production at the Tevatron, with this end in mind. The remarkable thing is that the non-perturbative parameters appearing in the  $\eta_c$  production cross section can be determined from the matrix elements determined from  $J/\psi$  production at the Tevatron: this happens because of the heavy-quark symmetry of the NRQCD Lagrangian. This has been exploited earlier in the context of  $h_c$  production at Tevatron [14]. We show, in the following, that from NRQCD one expects a large cross section for  $\eta_c$  production at the Tevatron. The canonical search channel for the  $\eta_c$  is its two photon decay mode and the number of events is substantial even after taking the small branching ratio of the  $\eta_c \rightarrow \gamma\gamma$  into account.

By writing a Fock space expansion of the physical  $\eta_c$ , which is a  $^1S_0$  ( $J^{PC} = 0^{-+}$ ) state, we have

$$|\eta_c\rangle = \mathcal{O}(1) |Q\bar{Q}[^1S_0^{[1]}]\rangle + \mathcal{O}(v^2) |Q\bar{Q}[^1P_1^{[8]}]g\rangle + \mathcal{O}(v^4) |Q\bar{Q}[^3S_1^{[8]}]g\rangle + \dots \quad (2)$$

The colour-singlet  $^1S_0$  state contributes at  $\mathcal{O}(1)$  but the colour-octet  $^1P_1$  and  $^3S_1$  channels effectively contribute at the same order because the  $P$ -state production is itself down by factor of  $\mathcal{O}(v^2)$ . The colour-octet states become a physical  $\eta_c$  by the  $^1P_1^{[8]}$  state emitting a gluon in an E1 transition, while the  $^3S_1^{[8]}$  emitting a gluon in

an M1 transition. To  $\mathcal{O}(\alpha_s^3 v^7)$  the  $\eta_c$  production cross section is

$$\begin{aligned} \sigma(\eta_c) = & \frac{F_1[{}^1S_0]}{M^2} \langle 0 | \mathcal{O}_1^{\eta_c}[{}^1S_0] | 0 \rangle \\ & + \frac{F_8[{}^1P_1]}{M^4} \langle 0 | \mathcal{O}_8^{\eta_c}[{}^1P_1] | 0 \rangle + \frac{F_8[{}^3S_1]}{M^2} \langle 0 | \mathcal{O}_8^{\eta_c}[{}^3S_1] | 0 \rangle. \end{aligned} \quad (3)$$

The coefficients  $F$ 's are the cross sections for the production of  $c\bar{c}$  pair in the respective angular momentum and colour states and is given by

$$\begin{aligned} \frac{d\sigma}{dp_T} (p\bar{p} \rightarrow c\bar{c} [{}^{2S+1}L_J^{[1,8]}] X) = \\ \sum \int dy \int dx_1 \, x_1 \, G_{a/p}(x_1) \, G_{b/\bar{p}}(x_2) \frac{4p_T}{2x_1 - \bar{x}_T e^y} \frac{d\hat{\sigma}}{d\hat{t}}(ab \rightarrow c\bar{c} [{}^{2S+1}L_J^{[1,8]}] d). \end{aligned} \quad (4)$$

The contributing subprocess cross sections are

$$\begin{aligned} q \, \bar{q} & \rightarrow Q\bar{Q}[{}^{2S+1}L_J] \, g, \\ g \, q(\bar{q}) & \rightarrow Q\bar{Q}[{}^{2S+1}L_J] \, q(\bar{q}), \\ g \, g & \rightarrow Q\bar{Q}[{}^{2S+1}L_J] \, g, \end{aligned} \quad (5)$$

where the  $Q\bar{Q}$  is in the  ${}^1S_0^{[1]}$ ,  ${}^1P_1^{[8]}$  and  ${}^3S_1^{[8]}$  states. The summation is over the partons,  $a$  and  $b$ .  $G_{a/p}$  and  $G_{b/\bar{p}}$  are the distributions of partons  $a$  and  $b$  in the hadrons and  $x_1$  and  $x_2$  are the momentum they carry, respectively.  $x_2$  is given in terms of  $x_1$  as

$$x_2 = \frac{x_1 \bar{x}_T e^{-y} - 2\tau}{2x_1 - \bar{x}_T e^y}, \quad (6)$$

where  $\tau = M^2/s$ , with  $M$  the mass of the resonance,  $\sqrt{s}$  the centre-of-mass energy,  $y$  the rapidity at which the resonance is produced and  $\bar{x}_T = \sqrt{x_T^2 + 4\tau} \equiv 2M_T/\sqrt{s}$ , with  $x_T = 2p_T/\sqrt{s}$ .

Among the coefficients the matrix elements for the subprocesses corresponding to  $F_1[{}^1S_0]$  are available in [15] and those corresponding to  $F_8[{}^3S_1]$  are calculated by Cho and Leibovich [7]. We have calculated the remaining coefficient  $F_8[{}^1P_1]$ . The packages FORM and MATHEMATICA are used to calculate the matrix element squares of the subprocesses.

One important component at large- $p_T$  is the gluon fragmentation to charmonium, as has been shown in the case of large- $p_T$   $J/\psi$  production [4]. It has been noted in Ref. [6] that even in the fixed-order calculation of the  $J/\psi$  cross section, the diagrams that contribute at large- $p_T$  are fragmentation-like diagrams in which a single gluon attaches itself to a  $c\bar{c}$  pair. In our case, we find that while the  $F_8[{}^3S_1]$  has

a fragmentation-like contribution at large- $p_T$  there is no such contribution for the  $F_8[{}^1P_1]$ . Following Ref. [6], we have also corrected the cross section by the ratio of  $Q^2$  evolved to the unevolved fragmentation function to get the correct shape of the  $p_T$  distribution at large  $p_T$ .

Heavy quark spin-symmetry is made use of in obtaining  $\langle \mathcal{O}_n^{\eta_c} \rangle$ 's from the experimentally available  $\langle \mathcal{O}_n^{J/\psi} \rangle$ 's. Using this symmetry we get the following relations among  $\langle \mathcal{O}_n^H \rangle$ 's:

$$\begin{aligned} \langle 0 | \mathcal{O}_1^{\eta_c} [{}^1S_0] | 0 \rangle &= \langle 0 | \mathcal{O}_1^{J/\psi} [{}^3S_1] | 0 \rangle (1 + O(v^2)), \\ \langle 0 | \mathcal{O}_8^{\eta_c} [{}^1P_1] | 0 \rangle &= \langle 0 | \mathcal{O}_8^{J/\psi} [{}^3P_0] | 0 \rangle (1 + O(v^2)), \\ \langle 0 | \mathcal{O}_8^{\eta_c} [{}^3S_1] | 0 \rangle &= \langle 0 | \mathcal{O}_8^{J/\psi} [{}^1S_0] | 0 \rangle (1 + O(v^2)). \end{aligned} \quad (7)$$

Hence, for the singlet matrix elements we have  $\langle 0 | \mathcal{O}_1^{J/\psi} [{}^3S_1] | 0 \rangle = 1.2 \text{ GeV}^3$  [6] and for the octet matrix elements extracted from the CDF  $J/\psi$  data we have  $A_1 + A_2 \equiv \frac{\langle 0 | \mathcal{O}_8^{J/\psi} [{}^3P_0] | 0 \rangle}{M_c^2} + \frac{\langle 0 | \mathcal{O}_8^{J/\psi} [{}^1S_0] | 0 \rangle}{3} = (2.2 \pm 0.5) \times 10^{-2} \text{ GeV}^3$  [7]. The CDF  $J/\psi$  do not allow for a separate determination of the values of  $A_1$  and  $A_2$  because the shapes of these two contributions to the  $J/\psi$   $p_T$  distribution are almost identical. We use the fact that the sum  $A_1 + A_2$  is constrained by the CDF data, and for our numerical predictions we assume that either  $A_1$  or  $A_2$  saturates the sum. In one case, therefore, we have the maximum possible contribution from the  ${}^3S_1^{[8]}$  channel and none from the  ${}^1P_1^{[8]}$  channel, and in the other case the  ${}^3S_1^{[8]}$  channel makes no contribution while the  ${}^1P_1^{[8]}$  contributes its maximum.

In Fig. 1, we have plotted the differential cross section  $d\sigma/dp_T$  against  $p_T$ . We are considering the case where the  $\eta_c$  will be searched for in its decay into two photons and we have consequently folded in the differential cross section with the  $\eta_c \rightarrow \gamma\gamma$  branching ratio ( $B = 3 \times 10^{-4}$ ). In our computations, we have used MRSD-' parton densities [16] and the parton densities are evolved to a scale  $Q = M_T$ . As a consequence of the fact that we are saturating the sum with either  $A_1$  or  $A_2$ , our curves for the two octet channels represent the maximum possible cross section that can result in each (with the other being zero). Because of the fact that fragmentation dominates at high  $p_T$  contributions from  ${}^1S_0^{[1]}$  and  ${}^1P_1^{[8]}$ , which do not have a fragmentation contribution, fall steeply with increasing  $p_T$ , whereas the  ${}^3S_1^{[8]}$  contribution makes a sizable contribution even at the largest  $p_T$  considered. The shape of the measured  $p_T$  distribution is clearly a very good measure of the individual matrix elements  $A_1$  and  $A_2$ . We reiterate that the CDF  $J/\psi$  data does not offer such a discrimination.

To get an idea of the measurability of the  $\eta_c$  cross section at the Tevatron, we have integrated our  $p_T$  distribution for values of  $p_T$  above 5 GeV. Assuming an integrated luminosity of  $110 \text{ pb}^{-1}$  we find that the number of events in the  $\gamma\gamma$  channel from the singlet  ${}^1S_0^{[1]}$  is about 400 and that from  ${}^3S_1^{[8]}$  is about 7300 when  ${}^1P_1^{[8]}$  contribution




Figure 1:  $d\sigma/dp_T$  (in nb/GeV) for  $\eta_c$  production (after folding in with  $\text{Br}(\eta_c \rightarrow \gamma\gamma) = 3.0 \times 10^{-4}$ ) in  $p\bar{p}$  collisions at 1.8 TeV with  $-0.6 \leq y \leq 0.6$ .

is absent, while the number of events from  $^1P_1^{[8]}$  is about 25 when  $^3S_1^{[8]}$  contribution is absent. Thus the number of  $\eta_c \rightarrow \gamma\gamma$  lies between 425 and 7700. Thus, even the minimum of 425 events provides a very good prospect for the observation of  $\eta_c$  at the Tevatron. We also point out that the integrated cross sections are also good measures of the matrix elements  $A_1$  and  $A_2$ .

We have studied the effect on the cross section of the variation of the parton densities, the scale and the non-perturbative matrix elements. By using GRV HO densities [17], instead of MRSD-', we find that the cross section increases by about 25%. If we use the scale choice  $Q = M_T/2$  instead of  $Q = M_T$  then the cross section increases by about 45-50%. In addition to these uncertainties, we can expect a 30% variation in the values of the non-perturbative matrix elements we have used, because of the fact that the heavy-quark symmetry is approximate.

In our calculations, we have considered only the direct production of  $\eta_c$ . There is a contribution to the  $\eta_c$  signal coming from the decays of  $J/\psi$ 's. From the measured

$J/\psi$  cross section at CDF we find that the contribution to the signal coming from  $J/\psi$  decays will be roughly about 100 events which is only a small fraction of the minimum number of events that we expect from the direct production process. A more complete analysis including the  $J/\psi$  decay contribution to the  $\eta_c$  signal will be presented in a future publication.

We conclude with the following observations: the heavy-quark symmetry of the NRQCD Lagrangian allows us to make predictions for the  $\eta_c$  cross section at the Tevatron. The integrated rate and the shape of the  $p_T$  distributions are very sensitive to the non-perturbative matrix elements  $A_1$  and  $A_2$ , whose sum has been determined from the  $J/\psi$  production data from CDF, but are not known individually. This sensitivity can clearly be used to determine these matrix elements individually. Our estimates for the integrated rate vary between 425 and 7700  $\eta_c$  events (after including the branching ratio of the  $\eta_c$  into two photons). Even at its minimum of 425 events, the prospect of observability of the  $\eta_c$  signal at the Tevatron seems extremely good. We emphasise that this is a testable prediction of NRQCD, and that such a prediction is not possible in alternate approaches to quarkonium formation like the colour evaporation model.

## References

- [1] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev. D* **46** R1914 (1992).
- [2] F. Abe *et. al*, *Phys. Rev. Lett.* **71**, 2537 (1993).
- [3] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev. D* **51**, 1125 (1995);erratum *ibid.* **55** 5853 (1997).
- [4] E. Braaten, M.A. Doncheski, S. Fleming and M. Mangano, *Phys. Lett. B* **333** 548 (1994); D.P. Roy and K. Sridhar, *Phys. Lett. B* **339** 141 (1994); M. Cacciari and M. Greco, *Phys. Rev. Lett.* **73** 1586 (1994).
- [5] E. Braaten and S. Fleming, *Phys. Rev. Lett.* **74**, 3327 (1995).
- [6] P. Cho and A.K. Leibovich, *Phys. Rev. D* **53**, 150 (1996).
- [7] P. Cho and A.K. Leibovich, *Phys. Rev. D* **53**, 6203 (1996).
- [8] S. Aid *et. al.*, H1 Collaboration, *Nucl. Phys. B* **472**, 3 (1996); ZEUS Collaboration, presented at the XXVIII Conference on High Energy Physics, Warsaw, July 1996, Ref. Num. pa 02-47.
- [9] M. Cacciari and M. Kramer, *Phys. Rev. D* **76**, 4128 (1996).
- [10] B.A. Kniehl and G. Kramer, DESY Preprint DESY 98-023, hep-ph/9803256.
- [11] K. Sridhar, A.D. Martin and W.J. Stirling, To appear in *Phys. Lett. B*, hep-ph/9806253.
- [12] M. Beneke, I. Z. Rothstein and M. B. Wise, *Phys. Lett. B* **408**, 373 (1997).
- [13] P. Cho and M. Wise, *Phys. Lett. B* **346** 129 (1995); S. Baek, P. Ko, J. Lee and H.S. Song, *Phys.Rev. D* **55** 6839 (1997); M. Beneke and I. Rothstein, *Phys. Lett. B* **372** 157 (1996); M. Beneke and M. Krämer, *Phys. Rev. D* **55** 5269 (1997).
- [14] K. Sridhar, *Phys. Rev. Lett.*, **77**, 4880 (1996).
- [15] R. Gastmans, T. Troost and T.T. Wu, *Nucl. Phys. B* **291** 731 (1996).
- [16] A.D. Martin, R.G. Roberts and W.J. Stirling, *Phys. Lett. B* **306** 145 (1993); *Phys. Lett. B* **309** 492 (1993).
- [17] M. Glück, E. Reya and A. Vogt, *Phys. Lett. B* **306** 391 (1993).